

Geodynamics of lithosphere as one of the crucial factors of mineral deposits formation of Ukraine

© V. Starostenko, O. Gintov, R. Kutas, I. Pashkevich, 2010

Institute of Geophysics, National Academy of Sciences of Ukraine, Kiev, Ukraine

earth@igph.kiev.ua

kutro@ndc.org.ua

oleg.gintov@gmail.com; innap@34mail.ru

Comprehensive studies of the lithosphere structure of Ukraine have been performed lately at the Institute of Geophysics NASU in connection with solving the problems of metallogeny, diamond-, gas- and oil-bearing [Starostenko et al., 2007]. These studies included: thorough analysis of geological-geophysical data on tectonics and deep structure of the Earth's crust and upper mantle; plotting the comprehensive three-dimensional geophysical and geodynamic model of lithosphere; generalization of global experience in application of geophysical, including tectonophysical, methods for the search and prospecting of different kinds of mineral deposits. The results of studies allow drawing some conclusions on the character of geodynamic processes, having an influence on formation and distribution of deposits on the territory of Ukraine.

It has been shown by the examples of the Ukrainian Shield, the Dnieper-Donets, the Carpathian and Azov-Black Sea oil-gas-bearing provinces that the regularities of formation, redistribution and concentration of mineral deposits are determined in many cases by special features of tectonosphere evolution arising from the mechanisms of global and regional movements of lithosphere plates.

The Ukrainian Shield (USh). Practically all large fault zones of the USh are *zones of shift with acute predominance of horizontal component of banks displacements* [Gintov, 2005]. They were arisen in Neo-Archean and Early Proterozoic as right and left shifts, strike-slip and upthrow faults with amplitude of displacement as kilometers and tens of kilometers and transcending far beyond the limits of the Shield. Their roots according to different geophysical characteristics are related to in the mantle up to the depth of 100–200 km and more [Gintov, Pashkevich, 2010]. Existence of such ancient extensive zones of shift may be only explained by convective movements in the mantle, which took place as early as in Neo-Archean and Early Proterozoic. In Proterozoic numerous fault zones were

also formed as listric ones or gently sloping overthrust.

According to the results of tectonophysical and paleomagnetic studies of dynamics of lithosphere of the USh a conclusion has been drawn that the Shield as a consolidated structure has existed since the boundary of 1.8–1.6 Ga. According to paleomagnetic data [Elming et al., 1998], it existed before and moved autonomously and even earlier, according to tectonophysical data [Gintov, Pashkevich, 2010], it was divided into several megablocks, which had their own trajectories of movements.

Joint analysis of kinematic and three-dimensional geophysical model of the Shield as well as of V_p tomographic model of the mantle up to the depth of 850 km [Geyko et al., 2006], allows to correlate the results of metallogenic studies of fault zones of the USh with materials of geodynamic reconstructions. Within the limits of the USh more than 75 percent of known metallogenic zones, ore areas and ore fields belong to well studied large fault zones. These are mainly the areas of development of mineralization and deposits of non-ferrous, rare, noble metals, uranium, rare earths et al. [Starostenko et al., 2007].

Dnieper-Donets oil-gas province (DDOGP). Numerous oil-gas deposits within non-anticline, the so-called "non-traditional" traps — on monoclines, half-anticlines, sub-thrust zones, within crystalline basement etc. have been discovered here lately. The majority of such deposits are related to the fractures of fault-, shift and upthrow (shear) fault types. These considerations brought a lot of scientists to the idea of great perspectives of anorganic hypothesis of oil origin and of important role of tangential forces and horizontal movements of lithosphere in formation of traps for hydrocarbons.

Numerical modeling of the process of the Dnieper-Donets aulacogene (DDA) formation by the back-stripping method within the limits of continental lithosphere stretching concept of D. McKenzie testifies the possibility of its formation in the Late

Devonian by the type of rift basins with formation of sub-oceanic crust [Stifenson et al., 1997]. Important role has been proved of shifting processes in formation of contemporary structure of DDOG and of many types of oil-gas deposits. The system of faults of the DDA manifested during Alpine time as the largest right shift, which is the result of super-regional sub-longitudinal contraction, which covered during Meso-Cenozoic time the territories of the south and south-west of the East-European Platform (EEP). The explanation can be found in the known plate tectonic reconstructions: the pressure on the EEP from the south as a result of movements of the African and Arabian plates is passed through West-Black Sea and East Black Sea micro-plates [Nikishin et al., 2001; Patalaha et al., 2003; Kazmin et al., 2004].

The suture zone of the Donets Folded Structure (DFS) with the south slope of the Voronezh crystalline massif (Starobelsk-Millerov monocline) can be considered as an example of the influence of the processes of compression and shifting upon formation of hydrocarbon deposits. The suture zone is revealed by the series of over-thrusts of Carboniferous-Cretaceous deposits of the DFS upon the monocline — Krasnopopovskiy, Severodonetskiy, Mari-evskiy, Almaznyi, Iliychevskiy and other ones. Here, within the stripe not less than 50 km wide, a whole set of non-anticline type deposits within the limits of the monocline itself, in Krasnorechensk and Lisichansk gas-bearing areas have been discovered [Gintov, 2005; Starostenko et al., 2009].

The Carpathian Meso-Cenozoic Oil-Gas Province (COGP) is characterized by over-thrust structure of the Meso-Cenozoic strata, doubled and tripled cross-section of the Cretaceous, Paleogene and Neogene, is considered by the majority of geologists and geophysicists from the positions of plate tectonics.

According to the data of seismic tomography high-velocity lithosphere of EEP sinks under relatively low-velocity mantle of the Volyno-Podolian plate and the Carpathians from the depth of 50 to 250—300 km [Geyko et al., 2006].

It has been found by geothermic studies [Kutas, 2005] that for deposits and the areas of oil-gas accumulation, concentrated within the limits of Pre-Carpathian depression (in this case within the External zone of depression gas and gas-condensate deposits predominate, and within Internal one — oil deposits), increased temperature and heat flows are specific. Deposits form two stripes, narrow enough in zones of Carpatian direction Pre-Carpathian and Scole faults which are at a distance of 20—30 km from each other.

In the periphery of EEP accumulation of hydrocarbons in the deposits of accretion wedge could occur as early as Cretaceous and Paleogene. During collision stage redistribution of hydrocarbon potential occurred. Migration processes influenced essentially on hydrodynamic conditions, thermal regime, physical parameters of sedimentary strata. These parameters can be used for determination of the ways of migration and zoning of oil-gas areas.

With the help of numerical tectonophysical modeling the connection of formation and distribution of oil-gas deposits COGP with plate-tectonic mechanism of region formation has been analyzed [Gonchar, 2007]. Palinspastic sections of the frontal part of the Flish Carpathians and Pre-Carpathian depression have been completed. They reconstruct the process of formation of deposits under conditions of lateral accretion and formation of covers in case of obduction-subduction mechanism and explain zoning in distribution of oil and gas deposits within Pre-Carpathian depression.

Application of modern geodynamic concepts of the Carpathian region formation allows considerable widening of its perspectives of oil- and gas-bearing. North-western part of the External zone of Pre-Carpathian depression (Krukenitskaya sub-zone), autoktonous deposits of Folded Carpathians, platform Meso-Cenozoic deposits of south-eastern part of External zone under the over-thrust of the Folded Carpathians, Trans-Carpathian depression are worth to be paid special attention.

Azov-Black Sea Oil-Gas Province (ABSOGP). A considerable part of the Province is located within the water areas. Therefore analysis of geodynamic processes in it is possible only on the background of geodynamic development of the whole Black Sea-Caucasian segment of the Alpine belt.

ABSOGP and COGP have got some common geophysical peculiarities. It allows treating them from similar geodynamic positions. ABSOGP adjoins to the southern tectonic border of EEP and here, as well as in the COGP, according to the data of seismic tomography [Geyko et al., 2006] submergence can be observed of relatively high-velocity lithosphere of the Craton under relatively low-velocity mantle of Dobrogea, Scythian plate, the Black Sea. The studies of geothermal regime of the region shows that oil-gas deposits are localized within the areas of increased heat flows, forming in this case two sub-parallel stripes: gas deposits are located within the zones of high heat flows and the oil ones — in their periphery. The important role is played by tectono-thermal activation in case of shift displacements of micro-plates [Kutas et al., 2002; 2007].

For the studies of ABSOGP conditions of formation the main interest belongs to Oligocene-Quaternary stage, which began 35—40 million years ago. The results of geothermal studies in the region show that for this period there is strong enough correlation between location of oil-gas deposits and anomalies of heat flow [Kutas et al., 2002; 2007]. At this stage the Black Sea — Caucasian region develops in the indenter regime under the action of Arabian, Pannonian and Adriatic plates [Patalaha et al., 2003]. In addition, the movement of the Black Sea plate within northern points of compass was of reverse character that is indicated by tectonophysical data on the Crimean peninsula: the phases of

sub-longitudinal contraction was interrupted by shorter phases of even stretching [Gintov, 2005].

For promising areas of ABSOGP — north-western shelf of the Black Sea and the Kerch-Tamanian depression the detailed schemes of fault tectonics of consolidated crust have been compiled [An integrated ..., 2006]. They emphasize the leading role of shift deformations during formation of fault and fold structures of the Meso-Cenozoic cover, to which (especially to the knots of fault crossing) hydrocarbon deposits are related. It has also been found by marine geophysical studies that zones of deep faults and neo-tectonic disturbances connected with them are the channels of migration of gas-fluid flows [Kobolev, Kutas, 1999].

References

- An integrated* three-dimensional geophysical model of the lithosphere of the Ukrainian Shield in connection with magmatism, tectonics and the formation of minerals // Report of the Institute of Geophysics by S.I. Subbotin of NASU / Ed. V. I. Starostenko. — Kiev, 2006. — 510 p. — Fund of UkrNIINTI, № state registration 102U002478 (in Russian)
- Elming S. A., Mikhailova N. P., Kravchenko S. N.* The Nonconsolidation of the East European Craton; a Paleomagnetic Analysis of Proterozoic Rocks from the Ukrainian Shield and Tectonic reconstructions Versus Fennoscandia // *Geophys. J.* — 1998. — **20**, № 4. — P. 71—74 (in Russian).
- Geyko V. S., Shumlanskaya L. A., Bugaenko I. V., Zayets L. N., Tsvetkova T. A.* Three-dimensional model of the upper mantle of Ukraine on the arrival time of *P*-waves // *Geophys. J.* — 2006. — **28**, № 1. — P. 3—16 (in Russian).
- Gintov O. B.* Field tectonophysics and its application in studying the crustal deformation of Ukraine. — Kiev: Phoenix, 2005. — 572 p. (in Russian).
- Gintov O. B.* Tectonophysics in solving important economic problems (review of research in the CIS countries). 1 // *Geophys. J.* — 2009. — **31**, № 5. — P. 3—31 (in Russian).
- Gintov O. B., Pashkevich I. K.* Tectonophysical analysis and geodynamic interpretation of the three-dimensional geophysical model of the Ukrainian Shield // *Geophys. J.* — 2010. — **32**, № 2. — P. 3—27 (in Russian).
- Gonchar V. V.* Rheological control of the accretion and cover styles of deformation and stress state of suprasubduction orogen // *Geophys. J.* — 2007. — **29**, № 6. — P. 116—137 (in Russian).
- Kazmin V. G., Lobkovsky L. I., Pustovitenko B. G.* Contemporary kinematics of the microplates in the Black-sea-South-Caspian region // *Okeanology.* — 2004. — **44**, № 4. — P. 600—610 (in Russian).
- Kobolev V. P., Kutas R. I.* Geophysical researches of gas release structures on the northwestern part of the Black Sea // *Tectonics and oil- and gas-content of the Azov-Black-sea region.* — Simferopol, 1999. — P. 53—55 (in Russian).
- Kutas R. I.* Geodynamic processes and thermal state of lithosphere of the Carpathian region // *Research of modern geodynamics of the Ukrainian Carpathians.* — Kiev: Nauk. dumka, 2005. — P. 132—139 (in Russian).
- Kutas R. I., Kobolyev V. P., Paliy S. I.* Geothermal conditions and oil- and gas bearing of the North-Black-sea — Ciscaucasian region // *Oil and Gas Industry.* — 2002. — № 5. — P. 9—11 (in Russian).
- Kutas R. I., Kravchuk O. P., Bevzyuk M. I., Stakhova L. I.* Results of geothermal studies in the northern part of the Black Sea // *Geophys. J.* — 2007. — **29**, № 4. — P. 45—65 (in Russian).
- Nikishin A. M., Korotayev M. V., Bolotov S. N., Yershov A. V.* Tectonic history of the Black-sea basin // *BMOIP.* — 2001. — **76**, issue 3. — P. 3—17 (in Russian).
- Patalakha E. I., Gonchar V. V., Senchenkov I. K., Chervinko O. P.* Indenter mechanism in geodynamics of the Crimean-Black-sea region. Forecast of HC and seismic hazard. — Kiev: Publishing house PP "EMKO", 2003. — 226 p. (in Russian).
- Starostenko V. I., Gintov O. B., Pashkevich I. K., Burakhovich T. K., Kulik S. N., Kuprienko P. Ya., Makarenko I. B., Orlyuk M. I., Tsvetkova T. A.* Regularities in the distribution of ore mineral deposits in con-

nection with the deep structure and dynamics of the lithosphere of the Ukrainian Shield // *Geophys. J.* — 2007. — **29**, № 5. — P. 3—34 (in Russian).

Starostenko V. I., Lukin A. E., Kobolyev V. P., Ruskov O. M., Orlyuk M. I., Shuman V. N., Omelchenko V. D., Pashkevich I. K., Tolkunov A. P., Bogdanov Yu. A., Burkin'sky I. B., Loiko N. P., Fedotova I. N., Zakharov I. G., Chernyakov A. M., Kuprienko P. Ya, Makarenko I. B., Legostayeva O. V., Lebed T. V., Savchenko A. S. Model of deep struc-

ture of the Donets folded construction and surrounding structures according to the regional geophysical observations // *Geophys. J.* — 2009. — **31**, № 4. — P. 44—68 (in Russian).

Stephenson R. A., Van Weiss Ja. D., Stovba S. N., Shymanovsky V. A. One-dimensional numerical modeling of tectonic immersion of DD within the concept of extension of the continental lithosphere by Mac Kenzie // *Geophys. J.* — 1997. — **19**, № 3. — P. 25—41 (in Russian).