

? (17

... , ... , ... , ... , ... , ... , ... ,
... , ... , ... , ... , ... , ... , ... ,
... , ... , ... , ... (. - . . .),
«
5/7; « »,
».

OPTIMISM OR PESSIMISM? (17 YEARS OF FIGHTING PEDIATRIC HIV-INFECTION IN THE CRIMEA)

**N.V. Rymarenko, O.A. Zalata, G.L. Kiselyova, N.P. Kiselyova, I.G. Semenova, E.V. Dotsenko,
T.N. Bezdolnaya, A.E. Chitakova, E.R. azinova, L.I. Anohina, N.I. Yavorskaya, R.L. Belalov,
E.E. Karaeva, M.S. Ziyatdinova, E.N. Begma, N.A. Vasilyeva**

SUMMARY

The article presents an analysis of observation and treatment results for HIV-infected children in AR Crimea over the period of 17 years. There were 184 children under observation and 90% of them appeared to have been infected from the mother. After examination of 14 patients, we diagnosed the first clinical stage of the disease, in 47 (25%) – the second stage, in 65 (35%) – the third stage, and in 58 (32%) – the fourth stage of HIV infection (AIDS). The most often clinical signs observed in patients were enlarged peripheral lymphonodus (94%), regression of the weight (65%), and hepatosplenomegaly (9%). In total, during the whole period of observation, 14 (7%) children died due to the development of AIDS, and in 64% of cases the direct causes of death were pneumonia and tuberculosis. 169 children were taking antiretroviral therapy, and 15 (9%) were having it for more than 10 years. The article also addresses the main problems related to late diagnosis of HIV today and discusses possible solutions.

I I ?(17 I I -I)
... , ... , ... , ... , ... , ... , ... ,
... , ... , ... , ... , ... , ... , ... ,
... , ... , ... , ... , ... , ... , ... ,
17 . 184 , 90%
14 (8%) I , 47 (25%) – II , 65 (35%)
– III 58 (32%) – IV
(94%), (65%)
(9%). 14 (7%) ,
64% ,
169 , 15 (9%) – 10 .
:
187
[1].
: 1987 2013
9 413 / 17 , -
422,5/100
55,3/100

«
»
».
5, 12
5
:«
».
:
/
()
,
()
)

0-1%.

[1].

184 3 18
2007 , -
),
1-2
0,7% 2013 181 (98%)
), 165 (90%)
), 16 (8%) –
4-5), 2
, 1
-
CD4- (),
, 14 (8%)
I , 47
(25%) – II , 65 (35%) – III 58 (32%)
– IV ().
2000
2001 .,
1993 (76%)
, 22%
1996 , 2001 4
2% - 3
1
(n=184)

	.(%)
	173 (94%)
()	119 (65%)
	84 (46%)
-	17 (9%)
()	15 (8%)
-	5 (3%)
	5 (3%)
-	3 (2%)
:	
	93 (51%)
	21 (11%)
	22 (12%)

2014, 17, 2 (66) -

1- 132 (78%), 2- - 7 1. (4%) - 30 (18%) 3- - 7 (65%) (94%), (9%). 90% (51%)

(.2).

(n=184)

	(%)
	153 (84%)
	136 (74%)
	49 (27%)
	26 (14%)
	15 (8%)
	14 (7%)
	10 (5%)
	5 (2,5%)
	5 (2,5%)

(80%) - - 8%

2% - ()

(1%) 5% 22

(1%). 66% 8 3 (30%)

8%

()

zoster - 24% (herpes- labialis herpes- (SpO₂) 76-88%. 50%

14% 20% 20% -

(46%). 23% 10% - 7 (70%)

, 40% -
 , , -
 , , -
 1 2 (20%) , , -
 , 8 , , -
 , 8 , ! , -
 , , -
 , , -
 (, -
 -4 0 **09.04.2013** « **199** -
). » -
 , , -
 , 64% 14 (7%) , () . -
 , , - (-
 5 ,) , -
 , , -
 , , -
 , , -
 2013 . 2009- 2003-2004 . , -
 , 16 , , -
 2 3 , , 35 (83%) 42
 (13 18 , , -
) , , -
 6 2 , -
 , -4 , -
 5 (33%) 10 15 (38%) 35
 7 (27%) , -
 , 3 (20%) , 2003 , -
 3 (20%) , -
 , 3 (20%) , -
 , 7 (47%) , -
 , - , -
 5 , -
 (33%) 3 (20%) , -
 , 5 - , 35 (19%)
 (33%) 184, 18 2013 , -
 13 18 13 (38%) -
 , , -
 10 , -

