



ЗАМЕТКА

***Trichogramma cacoeiciae* Marchal, a New Species for the Fauna of England [*Trichogramma cacoeiciae* Marchal, новый вид для фауны Англии].** — Parthenogenetic strain of one *Trichogramma* species was reared from the eggs of *Mamestra brassicae* L. exposed on cabbage plants in the Chelsea Physic Garden (Center of London, England). In the French laboratory of INRA-INSA the strain was treated at hot temperature (30 degree all the cycle) for 7 generations. Only females were observed among the numerous individuals in each studied generation. So, the strain showed a thelytokous mode of reproduction which was not caused by the symbiotic bacteria of the genus *Wolbachia*. Only one species, *T. cacoeiciae* Marchal, is known to possess such a character. The esterases of the strain of *Trichogramma* were stained after the electrophoresis. All the 8 studied homogenates (20 females taken at random constituted one homogenate) were similar, with no polymorphism. Five bands (positions: 0.07, 0.24, 0.45, 0.48 and 0.51) corresponding to 4 alleles at 4 loci were identified: Est 1—0.07, Est 2—0.24, Est 5—0.45—0.48 and Est 6—0.51. It was shown that the esterase pattern of the strain from Chelsea corresponds to that of *T. cacoeiciae*. This determination confirms the diagnosis carried out from the absence of *Wolbachia*. This is the first discovery of *T. cacoeiciae* for the fauna of England. — V. N. Fursov (Schmalhausen Institute of Zoology, Kiev, Ukraine), B. Pintureau (Institut National de la Recherche Agronomique — INSA, Villeurbanne, France).

ЗАМЕТКА

The first record of the family Megamerinidae (Diptera, Acalyptrata) from the Ukraine [Первая находка семейства Megamerinidae (Diptera, Acalyptrata) в Украине]. — 2 ♀ of *Megamerina dolium* F. were collected in woods (Teremki, Kiev), 5—7.07.1998 (leg. L. Frantsevich & I. Hladun). — V. A. Korneyev (Schmalhausen Institute of Zoology, Kiev).

ЗАМЕТКА

The first record of the family Carnidae (Diptera, Acalyptrata) from the Ukraine [Первая находка семейства Carnidae (Diptera, Acalyptrata) в Украине]. — ♂ and 3 ♀ of *Hemeromyia anthracina* Collin were collected on a balcony window at my apartment (Lukyanyvka, Kiev), 12—13.05.1998 (leg. V. Korneyev). — V. A. Korneyev (Schmalhausen Institute of Zoology, Kiev).

ЗАМЕТКА

Коростель (*Crex crex* L.) на Казачьелагерской арене Нижнеднепровских песков [Landrail (*Crex crex* L.) on the Kazachelagerskaya arena of the Lower Dnieper Sands]. — Коростель не включен в Красную книгу Украины (1994), однако является одним из 14 видов птиц фауны нашей страны, занесенных в Европейский красный список. 1—5.06.1998 нами обследовалась западная, юго-западная и центральная части, 11—12.06.1998 — северная, северо-восточная и юго-восточная части Казачьелагерской арены, преимущественно в пределах Раденского военного полигона. Коростели отмечены в северной и северо-западной части необлесенной территории арены, где расположена низменность, вытянутая в направлении от с. Казачьи Лагери до пгт. Новая Маячка Цюрупинского р-на Херсонской обл. Обычно здесь распространены луга в комплексе с песчаной степью, но 1997—1998 гг. изобиловали осадками, поэтому в июне 1998 г. большая часть низменности была залита водой и преобладали мокрые луга и болота. 11 июня в двух точках северной части по голосу учтено по 2 коростеля, 12 июня в северо-западной части — еще один. Таким образом, в пределах данной низменности коростель является обычным гнездящимся видом. — Т. И. Котенко (Институт зоологии НАН Украины, Киев).