

1, 1, 2, 2
1, - ()-474002,
2, - ()-221005,
e-mail: naveengzp@gmail.com

NOSTOC MUSCORUM AG. EX BORN. ET

FLAH.

10 *Nostoc muscorum* Ag. ex Born. et Flah.
AKR 4
(), ()
)
10 6
1 2A

: , , *Nostoc muscorum*,
-
(Sharma et al., 2006, 2007).

(Sharma & Rai, 2006),

(Tormo et al., 2001).

(Fitzgeorge et al., 1994),

(Kirkpatrick et al., 2004).

(Woodcock, 1948),

« »

(Benson et al., 2005).

Karenia brevis

(Flewelling et al., 2005). (Turner et al., 1990),

10 18

« » *Microcystis aeruginosa* Kütz.,

;

« »

Nostoc

(Dodds et al., 1995).

(Vasconcelos et al., 1996). *Nostoc flagelliforme*,

(Gao, 1998).

“ ”, *N. rivulare*,

(Sivonen & Jones, 1999).

(*Nostoc* sp. strain 152; Sivonen et al., 1992; *Nostoc* sp. strain DUN901; Beattie et al., 1998; *Nostoc* sp. strain IO-102-I; Oksanen et al., 2004)

Nostoc muscorum

Ag. ex Born. et Flah.

(Cano et al., 1990; Bloor & England, 1991; El-Sheekh et al., 2006).

(Sharma et al., 2006)

N. muscorum

(Cheng et al., 2005),

9

Nostoc muscorum

Ag. ()

Nostoc muscorum

(*Nostocales*, *Nostocaceae*),

2,5 32

(Sharma et al., 2006). BG-11,

70 $\mu \cdot E \cdot m^{-2} \cdot s^{-1}$) / 14:10 (28 \pm 1 $^{\circ}C$.

(. . .) LD₅₀⁻¹ 14
(Krishnamurthy
et al., 1986),
1:4:15.
40 $^{\circ}C$.
(0,85 % NaCl);

, AKR, 6-8
25 \pm 2 $^{\circ}C$, 12:12,
ad libitum.

: (. . .)
(. . .), 36 , 18
18 : . . . , -
. . . .

(. . .)
(10, 20 30) ,
(0,85 % NaCl).
1 - μ L (w/v). 3-

(. . .) ()
(10, 20 30), (5, 10
15),

20
6

6 (Benson et al., 2005).
<20 +

5

()

7,4, 5 EDTA; 5 EGTA; 250 mM ; 50 2- NaCl).
(50 , pH
8
(; EC 3.1.3.16)
(1 000 x 10)

10 000 x 30
100 000 x 60

(Toivola et al., 1994).

P- P-
405 (An, Carmichael, 1994).

10


(Sharma et al., 2006).

(
(Tormo et al., 2001).

2006)

(Sharma, Rai,
N. muscorum.

(Gupta et al., 2003).



A – (600 \times); B – (600 \times).

N. muscorum

(Fitzgeorge et al., 1994) (Cheng et al., 2005). LD₅₀

(Gupta et al., 2003).

LR

16

(Benson et al., 2005)

(Slatkin et al., 1983).

1 2A,

(PP1 PP2) (

)

;

*N.K. Sharma*¹, *S. Singh*¹, *R. Bajpayi*² & *A.K. Rai*²

¹ SOS in Microbiology, Jiwaji University, Gwalior (MP) - 474002, India

² Department of Botany, Banaras Hindu University, Varanasi (UP) - 221005

e-mail: naveengzp@gmail.com

EFFECT OF *NOSTOC MUSCORUM* AG. EX BORN. ET FLAH. TOXINS ON UPPER
RESPIRATORY TRACT OF MICE

Intraperitoneal administration of 10 µL of *Nostoc muscorum* Ag. ex Born. et Flah. crude extract was sufficient enough for the death of male AKR strain mice at 4h of exposure. Body weight, behavioral and histological observations (necrosis of hepatic cells), and biochemical tests (protein phosphatase inhibition)

indicated for the presence of microcystines in the crude extract. Since, the cyanobacterium was airborne, the very doses were also administered intranasally into mice. A dose of 10 µL of the crude extract resulted in severe morphological alterations of nasal mucosa at 6h of exposures, but not beyond nasal cavity. Though, presence of protein phosphatases 1 and 2A in the tracheal tissues confirmed the susceptibility of the upper respiratory tissues against microcystins. However, due to the deposition of insufficient amount of toxins post-nasal cavity respiratory tissues exhibited no response. This indicated that the route of exposure plays an important role in conferring toxicity.

Keywords: airborne algae, hepatotoxic, inhalation, intraperitoneal, intranasal, *Nostoc muscorum*, protein phosphatases.

- *An, J.S & W.W. Carmichael. 1994. Use of a colorimetric protein phosphatase inhibition assay and enzyme linked immunosorbent assay for the study of microcystins and nodularins. *Toxicon* **32**: 1495-1507.
- Beattie, K.A., K. Kaya, T. Sano & G. Codd. 1998. Three dehydrobutyrine-containing microcystins from *Nostoc*. *Phytochemistry* **47**: 1289-1292.
- Benson, J.M., J.A. Hutt, K. Rein, S.E. Boggs, E.B. Barr & L.E. Fleming. 2005. The toxicity of microcystin-LR in mice following 7 days of inhalation exposure. *Toxicon* **45**: 691-698.
- Bloor, S. & R.R. England. 1991. Elucidation and optimization of the medium constituents controlling antibiotic production by the cyanobacterium *Nostoc muscorum*. *Enzyme Microbiol. Technol.* **13**: 76-81.
- Cano, M.M.S., M.C.Z. Mule, C.Z. Caire & D.R. Halperin. 1990. Inhibition of *Candida albicans* and *Staphylococcus aureus* by phenolic compounds from the terrestrial cyanobacterium *Nostoc muscorum*. *J. Appl. Phycol.* **2**: 79-82.
- Cheng, Y.S., T.A. Villareal, Y. Zhou et al. 2005. Characterization of red tide aerosol on the Texas coast. *Harmful Algae* **4**: 87-94.
- Dodds, W.K., D.A. Guddar & D. Mollenhauer. 1995. The ecology of *Nostoc*. *J. Phycol.* **31**: 2-18.
- El-Sheekh, M.M., M.E.H. Osman, M.A. Dyab & M.S. Amer. 2006. Production and characterization of antimicrobial active substances from the cyanobacterium *Nostoc muscorum*. *Environ. Toxicol. Pharm.* **21**: 42-50.
- Fitzgeorge, R.B., S.A. Clark & C.W. Keevil. 1994. Routes of intoxication. Pp. 66-74 in: *Detection methods for cyanobacterial toxins* / Eds. G.A. Codd., T.M. Jefferies., C.W. Keevil & C Potter. Roy. Soc. Chem., London.
- Flewelling, L.J., J.P. Narr & J.P. Abbott et al. 2005. Red tides and marine mammal mortalities. *Nature* **435**: 755-756.
- Gao, K. 1998. Chinese studies on the edible blue-green alga, *Nostoc flagelliforme*: a review. *J. Appl. Phycol.* **10**: 37-49.
- Gupta, N., S.C. Pant, P.V. Vijayaraghavan & P.V.L. Rao. 2003. Comparative toxicity evaluation of cyanobacterial cyclic peptide toxin microcystin variants (LR, RR, YR) in mice. *Toxicology* **188**: 285-296.
- Kirkpatrick, B., L.E. Fleming, D. Squicciarini et al. 2004. Literature review of Florida red tide: implications for human health effects. *Harmful Algae* **3**: 99-115.
- Krishnamurthy, T., W.W. Carmichael & E.W. Sarver. 1986. Toxic peptides from freshwater cyanobacteria (blue-green algae): Isolation, purification and characterization of peptides from *Microcystis aeruginosa* and *Anabaena flos-aquae*. *Toxicon* **24**: 865-873.

-
- Oksanen, I., J. Jokela, D.P. Fewer et al. 2004.** Discovery of rare and highly toxic microcystins from Lichen-associated cyanobacterium *Nostoc* sp. strain IO-102-I. *Appl. Environ. Microbiol.* **70**: 5756-5763.
- Sharma, N.K. & A.K. Rai. 2006.** Allergenicity of airborne cyanobacteria *Phormidium fragile* and *Nostoc muscorum*. *Ecotoxicol. Environ. Saf.* (DOI 10. 1016/j. ecoenv. 2006.08.06).
- Sharma, N.K., A.K. Rai, S. Singh & R.M.Jr. Brown. 2007.** Airborne algae: present status and their relevance. *J. Phycol.* **43**: 615-627.
- Sharma, N.K., S. Singh & A.K. Rai. 2006.** Diversity and seasonal variation in viable algal particles in a subtropical Indian City. *Environ. Res.* **102**: 252-259.
- Sivonen, K. & G. Jones. 1999.** Cyanobacterial toxins. Pp. 41-111 in: *Toxic cyanobacteria in water* / Eds. I. Chorus & J. Bartram. E & FN Spon, London.
- Sivonen, K., M. Namikoshi, W.R. Evans et al. 1992.** Three new microcystins, cyclic heptapeptide hepatotoxins, from *Nostoc* sp. strain 152. *Chem. Res. Toxicol.* **5**: 464-469.
- Slatkin, D.N., R.D. Stoner, W.H. Adams. 1983.** Atypical pulmonary thrombosis caused by toxic cyanobacterial peptide. *Science* **220**: 1383-1385.
- Toivola, D.M., J.E. Eriksson & D.L. Brautigam. 1994.** Identification of protein phosphatase 2A as the primary target for microcystin-LR in rat liver homogenates. *FEBS Lett.* **344**: 175-180.
- Tormo, R., D. Riccio, I. Silva & A.F. Muñoz. 2001.** A quantitative investigation of airborne algae and Lichen soredia obtained from pollen trap in South-West Spain. *Eur. J. Phycol.* **36**: 385-390.
- Turner, P.C., A.J. Gammie, K. Hollinrake & G.A. Codd. 1990.** Pneumonia associated with contact with cyanobacteria. *Brit. Med. J.* **300**: 1440-1441.
- Vasconcelos, V.M., K. Sivonen, W.R. Evans et al. 1996.** Hepatotoxic microcystin diversity in cyanobacterial blooms collected in Portuguese fresh water. *Water Res.* **30**: 2377-2385.
- Woodcock, A.H. 1948.** Note concerning human respiratory irritation associated with high concentration of plankton and mass mortality of marine organisms. *J. Mar. Res.* **7**: 56.

19.11.07